

# A NOTE ON THE FROBENIUS-EULER NUMBERS AND POLYNOMIALS ASSOCIATED WITH BERNSTEIN POLYNOMIALS

SERKAN ARACI AND MEHMET ACIKGOZ

**ABSTRACT.** The present paper deals with Bernstein polynomials and Frobenius-Euler numbers and polynomials. We apply the method of generating function and fermionic  $p$ -adic integral representation on  $\mathbb{Z}_p$ , which are exploited to derive further classes of Bernstein polynomials and Frobenius-Euler numbers and polynomials. To be more precise we summarize our results as follows, we obtain some combinatorial relations between Frobenius-Euler numbers and polynomials. Furthermore, we derive an integral representation of Bernstein polynomials of degree  $n$  on  $\mathbb{Z}_p$ . Also we deduce a fermionic  $p$ -adic integral representation of product Bernstein polynomials of different degrees  $n_1, n_2, \dots$  on  $\mathbb{Z}_p$  and show that it can be written with Frobenius-Euler numbers which yields a deeper insight into the effectiveness of this type of generalizations. Our applications possess a number of interesting properties which we state in this paper

## 1. Introduction and Notations

Let  $p$  be a fixed odd prime number. Throughout this paper we use the following notations. By  $\mathbb{Z}_p$  we denote the ring of  $p$ -adic rational integers,  $\mathbb{Q}$  denotes the field of rational numbers,  $\mathbb{Q}_p$  denotes the field of  $p$ -adic rational numbers, and  $\mathbb{C}_p$  denotes the completion of algebraic closure of  $\mathbb{Q}_p$ . Let  $\mathbb{N}$  be the set of natural numbers and  $\mathbb{N}^* = \mathbb{N} \cup \{0\}$ . The  $p$ -adic absolute value is defined by

$$|p|_p = \frac{1}{p}.$$

In this paper, we assume  $|q - 1|_p < 1$  as an indeterminate. In [17-19], let  $UD(\mathbb{Z}_p)$  be the space of uniformly differentiable functions on  $\mathbb{Z}_p$ . For  $f \in UD(\mathbb{Z}_p)$ , the fermionic  $p$ -adic integral on  $\mathbb{Z}_p$  is defined by T. Kim:

$$(1.1) \quad I_{-1}(f) = \int_{\mathbb{Z}_p} f(\xi) d\mu_{-1}(\xi) = \lim_{N \rightarrow \infty} \sum_{\xi=0}^{p^N-1} f(\xi) (-1)^\xi.$$

From (1.1), we have well known the following equality:

$$(1.2) \quad I_{-1}(f_1) + I_{-1}(f) = 2f(0)$$

here  $f_1(x) := f(x+1)$  (for details, see[3-24]).

---

2000 *Mathematics Subject Classification.* 05A10, 11B65, 28B99, 11B68, 11B73.

*Key words and phrases.* Frobenius-Euler numbers and polynomials, Bernstein polynomials, fermionic  $p$ -adic integral on  $\mathbb{Z}_p$ .

Let  $C([0, 1])$  be the space of continuous functions on  $[0, 1]$ . For  $C([0, 1])$ , the Bernstein operator for  $f$  is defined by

$$B_n(f, x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}$$

where  $n, k \in \mathbb{Z}_+ := \{0, 1, 2, 3, \dots\}$ . Here  $B_{k,n}(x)$  is called Bernstein polynomials, which are defined by

$$(1.3) \quad B_{k,n}(x) = \binom{n}{k} x^k (1-x)^{n-k}, \quad x \in [0, 1]$$

(for more informations on this subject, see [1-6, 11, 14, 15, 17, 21-24])

In [7], as is well known, Frobenius-Euler polynomials are defined by means of the following generating function:

$$(1.4) \quad \sum_{n=0}^{\infty} H_n(u, x) \frac{t^n}{n!} = e^{H(u, x)t} = \frac{1-u}{e^t - u} e^{xt}.$$

where the usual convention about replacing  $H^n(u, x)$  by  $H_n(u, x)$ . For  $x = 0$  in (1.4), we have to  $H_n(u, 0) := H_n(u)$ , which is called Frobenius-Euler numbers. Then, we can write the following

$$(1.5) \quad e^{H(u)t} = \sum_{n=0}^{\infty} H_n(u) \frac{t^n}{n!} = \frac{1-u}{e^t - u}.$$

By (1.4) and (1.5), we easily see the following applications:

$$\begin{aligned} e^{(H(u)+1)t} - ue^{H(u)t} &= 1 - u \\ \sum_{n=0}^{\infty} [(H(u) + 1)^n - uH_n(u)] \frac{t^n}{n!} &= 1 - u \end{aligned}$$

After these applications, we derive the following Lemma.

**Lemma 1.** For  $|u| > 1$  and  $n \in \mathbb{Z}_+ := \mathbb{N} \cup \{0\}$ , we have

$$(1.6) \quad (H(u) + 1)^n - uH_n(u) = \begin{cases} 1 - u, & \text{if } n = 0 \\ 0, & \text{if } n \neq 0. \end{cases}$$

In this paper, we obtained some relations between the Frobenius-Euler numbers and polynomials and the Bernstein polynomials. From these relations, we derive some interesting identities on the Frobenius-Euler numbers.

## 2. On the Frobenius-Euler numbers and polynomials

Let us take  $f(x) = u^x e^{tx}$  in (1.1), by (1.2), we see that

$$(2.1) \quad \int_{\mathbb{Z}_p} u^\eta e^{\eta t} d\mu_{-1}(\eta) = \frac{2}{1+u} H_n(-u^{-1}).$$

By (1.4) and (2.1), we have the following theorem.

**Theorem 1.**

$$(2.2) \quad \int_{\mathbb{Z}_p} u^\eta (x + \eta)^n d\mu_{-1}(\eta) = \frac{2}{u+1} H_n(-u^{-1}, x).$$

By applying some combinatorial techniques in (2.2), we derive the following

$$\int_{\mathbb{Z}_p} u^\eta (x + \eta)^n d\mu_{-1}(\eta) = \sum_{k=0}^n \binom{n}{k} x^{n-k} \left\{ \int_{\mathbb{Z}_p} u^\eta \eta^k d\mu_{-1}(\eta) \right\}.$$

So, from above, we have the well known identity

$$(2.3) \quad H_n(-u^{-1}, x) = \sum_{k=0}^n \binom{n}{k} x^{n-k} H_k(-u^{-1}) = (H(-u^{-1}) + x)^n.$$

by using the *umbral*(symbolic) convention  $H^n(u) := H_n(u)$ .

The Frobenius-Euler polynomials have to symmetric properties, which is shown by Choi *et al.* in [7], as follows:

$$H_n(-u^{-1}, 1-x) = (-1)^n H_n(-u^{-1}, x).$$

For  $n \in \mathbb{N}$ , by (2.3), Choi *et al.* derived the following equality:

$$(2.4) \quad u^2 H_n(-u^{-1}, 2) = u^2 + u + H_n(-u^{-1}).$$

From (2.2) and (2.4), we easily see that

$$(2.5) \quad \begin{aligned} & \int_{\mathbb{Z}_p} u^\eta (1-\eta)^n d\mu_{-1}(\eta) \\ &= (-1)^n \int_{\mathbb{Z}_p} u^\eta (\eta-1)^n d\mu_{-1}(\eta) \\ &= \frac{2}{u+1} (-1)^n H_n(-u^{-1}, -1) \\ &= \frac{2}{u+1} H_n(-u^{-1}, 2). \end{aligned}$$

Thus, we obtain the following Theorem.

**Theorem 2.** *The following identity*

$$(2.6) \quad \int_{\mathbb{Z}_p} u^\eta (1-\eta)^n d\mu_{-1}(\eta) = \frac{2}{u+1} H_n(-u^{-1}, 2)$$

*is true.*

Let  $n \in \mathbb{N}$ . By expression of (2.4) and (2.6), we get

$$(2.7) \quad \int_{\mathbb{Z}_p} u^\eta (1-\eta)^n d\mu_{-1}(\eta) = \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_n(-u^{-1}).$$

From (2.7), we procure the following corollary.

**Corollary 1.** *For  $n \in \mathbb{N}$ , we have*

$$\int_{\mathbb{Z}_p} u^\eta (1-\eta)^n d\mu_{-1}(\eta) = \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_n(-u^{-1}).$$

### 3. Some identities on the Frobenius-Euler numbers

In this section, we develop Frobenius-Euler numbers, that is, we derive some interesting and worthwhile relations for studying in Theory of Analytic Numbers.

Now also, for  $x \in [0, 1]$ , we rewrite definition of Bernstein polynomials as follows:

$$(3.1) \quad B_{k,n}(x) = \binom{n}{k} x^k (1-x)^{n-k}, \text{ where } n, k \in \mathbb{Z}_+.$$

By expression of (3.1), we have the properties of symmetry of Bernstein polynomials as follows:

$$(3.2) \quad B_{k,n}(x) = B_{n-k,n}(1-x), \text{ (for detail, see [21]).}$$

Thus, from Corollary 1, (3.1) and (3.2), we see that

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(\eta) u^\eta d\mu_{-1}(\eta) &= \int_{\mathbb{Z}_p} B_{n-k,n}(1-\eta) u^\eta d\mu_{-1}(\eta) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \int_{\mathbb{Z}_p} u^\eta (1-\eta)^{n-l} d\mu_{-1}(\eta) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \left( \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n-l}(-u^{-1}) \right). \end{aligned}$$

For  $n, k \in \mathbb{Z}_+$  with  $n > k$ , we compute

$$\begin{aligned} (3.3) \quad & \int_{\mathbb{Z}_p} B_{k,n}(\eta) u^\eta d\mu_{-1}(\eta) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \left( \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n-l}(-u^{-1}) \right) \\ &= \begin{cases} \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_n(-u^{-1}), & \text{if } k = 0, \\ \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \left( \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n-l}(-u^{-1}) \right), & \text{if } k > 0. \end{cases} \end{aligned}$$

Let us take the fermionic  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  on the Bernstein polynomials of degree  $n$  as follows:

$$\begin{aligned} (3.4) \quad \int_{\mathbb{Z}_p} B_{k,n}(\eta) u^\eta d\mu_{-1}(\eta) &= \binom{n}{k} \int_{\mathbb{Z}_p} \eta^k (1-\eta)^{n-k} u^\eta d\mu_{-1}(\eta) \\ &= \frac{2}{u+1} \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l H_{l+k}(-u^{-1}). \end{aligned}$$

Consequently, by expression of (3.3) and (3.4), we state the following Theorem:

**Theorem 3.** *The following identity holds true:*

$$\sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l H_{l+k}(-u^{-1}) = \begin{cases} 1 + u^{-1} + u^{-2} H_n(-u^{-1}), & \text{if } k = 0, \\ \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} (1 + u^{-1} + u^{-2} H_{n-l}(-u^{-1})), & \text{if } k > 0. \end{cases}$$

Let  $n_1, n_2, k \in \mathbb{Z}_+$  with  $n_1 + n_2 > 2k$ . Then, we derive the followings

$$\begin{aligned}
& \int_{\mathbb{Z}_p} B_{k,n_1}(\eta) B_{k,n_2}(\eta) u^\eta d\mu_{-1}(\eta) \\
&= \binom{n_1}{k} \binom{n_2}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{2k+l} \int_{\mathbb{Z}_p} (1-\eta)^{n_1+n_2-l} u^\eta d\mu_{-1}(\eta) \\
&= \left( \binom{n_1}{k} \binom{n_2}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{2k+l} \left( \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2-l}(-u^{-1}) \right) \right) \\
&= \begin{cases} \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2}(-u^{-1}), & \text{if } k=0, \\ \binom{n_1}{k} \binom{n_2}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{2k+l} \left( \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2-l}(-u^{-1}) \right), & \text{if } k \neq 0. \end{cases}
\end{aligned}$$

Therefore, we obtain the following Theorem:

**Theorem 4.** For  $n_1, n_2, k \in \mathbb{Z}_+$  with  $n_1 + n_2 > 2k$ , we have

$$\begin{aligned}
& \int_{\mathbb{Z}_p} B_{k,n_1}(\eta) B_{k,n_2}(\eta) u^\eta d\mu_{-1}(\eta) \\
&= \begin{cases} \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2}(-u^{-1}), & \text{if } k=0, \\ \binom{n_1}{k} \binom{n_2}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{2k+l} \left( \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2-l}(-u^{-1}) \right), & \text{if } k \neq 0. \end{cases}
\end{aligned}$$

By using the binomial theorem, we can derive the following equation.

$$\begin{aligned}
(3.5) \quad & \int_{\mathbb{Z}_p} B_{k,n_1}(\eta) B_{k,n_2}(\eta) u^\eta d\mu_{-1}(\eta) \\
&= \prod_{i=1}^2 \binom{n_i}{k} \sum_{l=0}^{n_1+n_2-2k} \binom{n_1+n_2-2k}{l} (-1)^l \int_{\mathbb{Z}_p} \eta^{2k+l} u^\eta d\mu_{-1}(\eta) \\
&= \frac{2}{u+1} \prod_{i=1}^2 \binom{n_i}{k} \sum_{l=0}^{n_1+n_2-2k} \binom{n_1+n_2-2k}{l} (-1)^l H_{2k+l}(-u^{-1}).
\end{aligned}$$

Thus, we can obtain the following Corollary:

**Corollary 2.** For  $n_1, n_2, k \in \mathbb{Z}_+$  with  $n_1 + n_2 > 2k$ , we have

$$\begin{aligned}
& \sum_{l=0}^{n_1+n_2-2k} \binom{n_1+n_2-2k}{l} (-1)^l H_{2k+l}(-u^{-1}) \\
&= \begin{cases} 1 + u^{-1} + u^{-2} H_{n_1+n_2}(-u^{-1}), & \text{if } k=0, \\ \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{2k+l} (1 + u^{-1} + u^{-2} H_{n_1+n_2-l}(-u^{-1})), & \text{if } k \neq 0. \end{cases}
\end{aligned}$$

For  $\eta \in \mathbb{Z}_p$  and  $s \in \mathbb{N}$  with  $s \geq 2$ , let  $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$  with  $\sum_{l=1}^s n_l > sk$ . Then we take the fermionic  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  for the Bernstein polynomials

of degree  $n$  as follows:

$$\begin{aligned}
& \int_{\mathbb{Z}_p} \underbrace{B_{k,n_1}(\eta) B_{k,n_2}(\eta) \dots B_{k,n_s}(\eta)}_{s\text{-times}} u^\eta d\mu_{-1}(\eta) \\
&= \prod_{i=1}^s \binom{n_i}{k} \int_{\mathbb{Z}_p} \eta^{sk} (1-\eta)^{n_1+n_2+\dots+n_s-sk} u^\eta d\mu_{-1}(\eta) \\
&= \prod_{i=1}^s \binom{n_i}{k} \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{l+sk} \int_{\mathbb{Z}_p} (1-\xi)^{n_1+n_2+\dots+n_s-l} u^\eta d\mu_{-1}(\eta) \\
&= \begin{cases} \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2+\dots+n_s}(-u^{-1}), & \text{if } k=0, \\ \prod_{i=1}^s \binom{n_i}{k} \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} \left( \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2+\dots+n_s-l}(-u^{-1}) \right), & \text{if } k \neq 0. \end{cases}
\end{aligned}$$

So from above, we have the following Theorem:

**Theorem 5.** For  $s \in \mathbb{N}$  with  $s \geq 2$ , let  $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$  with  $\sum_{l=1}^s n_l > sk$ . Then we have

$$\begin{aligned}
& \int_{\mathbb{Z}_p} u^\eta \prod_{i=1}^s B_{k,n_i}(\eta) u^\eta d\mu_{-1}(\eta) \\
&= \begin{cases} \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2+\dots+n_s}(-u^{-1}), & \text{if } k=0, \\ \prod_{i=1}^s \binom{n_i}{k} \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} \left( \frac{2}{u+1} + \frac{2}{u^2+u} + \frac{2}{u^3+u} H_{n_1+n_2+\dots+n_s-l}(-u^{-1}) \right), & \text{if } k \neq 0. \end{cases}
\end{aligned}$$

From the definition of Bernstein polynomials and the binomial theorem, we easily get

$$\begin{aligned}
& \int_{\mathbb{Z}_p} \underbrace{B_{k,n_1}(\eta) B_{k,n_2}(\eta) \dots B_{k,n_s}(\eta)}_{s\text{-times}} u^\eta d\mu_{-1}(\eta) \\
&= \prod_{i=1}^s \binom{n_i}{k} \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{\sum_{d=1}^s (n_d - k)}{l} (-1)^l \int_{\mathbb{Z}_p} \xi^{sk+l} u^\eta d\mu_{-1}(\eta) \\
(3.6) \quad &= \frac{2}{u+1} \prod_{i=1}^s \binom{n_i}{k} \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{\sum_{d=1}^s (n_d - k)}{l} (-1)^l H_{sk+l}(-u^{-1}).
\end{aligned}$$

Therefore, by (3.6), we get novel properties of Frobenius-Euler numbers with the following corollary:

**Corollary 3.** For  $s \in \mathbb{N}$  with  $s \geq 2$ , let  $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$  with  $\sum_{l=1}^s n_l > sk$ . Then, we have

$$\begin{aligned}
& u^2 \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{\sum_{d=1}^s (n_d - k)}{l} (-1)^l H_{sk+l}(-u^{-1}) \\
&= \begin{cases} u^2 + u + H_{n_1+n_2+\dots+n_s}(-u^{-1}), & \text{if } k=0, \\ \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} (u^2 + u + H_{n_1+n_2+\dots+n_s-l}(-u^{-1})), & \text{if } k \neq 0. \end{cases}
\end{aligned}$$

## REFERENCES

- [1] Açıkgöz, M. and Araci, S., A study on the integral of the product of several type Bernstein polynomials, IST Transaction of Applied Mathematics-Modelling and Simulation, vol.1, no. 1, pp. 10–14, 2010.
- [2] Açıkgöz, M. and Şimşek, Y., A New generating function of  $q$ -Bernstein type polynomials and their interpolation function, Abstract and Applied Analysis, Article ID 769095, 12 pages, doi: 10.1155/2010/769095.01-313.
- [3] Araci, S., Erdal, D., and Seo, J.-J., A study on the Fermionic  $p$ -adic  $q$ -integral Representation on  $\mathbb{Z}_p$  Associated with Weighted  $q$ -Bernstein and  $q$ -Genocchi Polynomials, Abstract and Applied Analysis, Volume 2011, Article ID 649248, 10 pages.
- [4] Araci, S. Erdal, D. and Kang, D.-J., Some New Properties on the  $q$ -Genocchi numbers and Polynomials associated with  $q$ -Bernstein polynomials, Honam Mathematical J. **33** (2011) no. 2, pp. 261-270
- [5] Araci, S., Acikgoz, M., Qi, F., On the  $q$ -Genocchi numbers and polynomials with weight 0 and their applications, <http://arxiv.org/abs/1202.2643>.
- [6] A. Bayad, T. Kim, Identities involving values of Bernstein  $q$ -Bernoulli, and  $q$ -Euler polynomials, Russ. J. Math. Phys. 18 (2011), no. 2, 133-143.
- [7] J. Choi, D. S. Kim, T. Kim and Y. H. Kim, A note on Some identities of Frobeniu-Euler Numbers and Polynomials, International Journal of Mathematics and Mathematical Sciences, Volume 2012, Article ID 861797, 9 pages.
- [8] T. Kim and B. Lee, Some Identities of the Frobenius-Euler polynomials, Abstract and Applied Analysis, Volume 2009, Article ID 639439, 7 pages.
- [9] T. Kim, On the multiple  $q$ -Genocchi and Euler numbers, Russian J. Math. Phys. 15 (4) (2008) 481-486. arXiv:0801.0978v1 [math.NT]
- [10] T. Kim, A Note on the  $q$ -Genocchi Numbers and Polynomials, Journal of Inequalities and Applications 2007 (2007) doi:10.1155/2007/71452. Article ID 71452, 8 pages.
- [11] T. Kim, A note  $q$ -Bernstein polynomials, Russ. J. Math. Phys. 18 (2011), 41-50.
- [12] T. Kim,  $q$ -Volkenborn integration, Russ. J. Math. Phys. 9 (2002), 288-299.
- [13] T. Kim,  $q$ -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients, Russ. J. Math. Phys. 15 (2008) , 51-57.
- [14] T. Kim, J. Choi, Y. H. Kim and C. S. Ryoo, On the fermionic  $p$ -adic integral representation of Bernstein polynomials associated with Euler numbers and polynomials, J. Inequal. Appl. 2010 (2010), Art ID 864247, 12pp.
- [15] T. Kim, J. Choi and Y. H. Kim Some identities on the  $q$ -Bernstein polynomials,  $q$ -Stirling numbers and  $q$ -Bernoulli numbers, Adv. Stud. Contemp. Math. 20 (2010) , 335-341.
- [16] T. Kim, An invariant  $p$ -adic  $q$ -integrals on  $\mathbb{Z}_p$ , Applied Mathematics Letters, vol. 21, pp. 105-108, 2008.
- [17] T. Kim, J. Choi and Y. H. Kim  $q$ -Bernstein Polynomials Associated with  $q$ -Stirling Numbers and Carlitz's  $q$ -Bernoulli Numbers, Abstract and Applied Analysis, Article ID 150975, 11 pages, doi:10.1155/2010/150975.
- [18] T. Kim,  $q$ -Euler numbers and polynomials associated with  $p$ -adic  $q$ -integrals, J. Nonlinear Math. Phys., 14 (2007), no. 1, 15–27.
- [19] T. Kim, New approach to  $q$ -Euler polynomials of higher order, Russ. J. Math. Phys., 17 (2010), no. 2, 218–225.
- [20] T. Kim, Some identities on the  $q$ -Euler polynomials of higher order and  $q$ -Stirling numbers by the fermionic  $p$ -adic integral on  $\mathbb{Z}_p$ , Russ. J. Math. Phys., 16 (2009), no.4, 484–491.
- [21] T. Kim, A. Bayad, Y. H. Kim, A Study on the  $p$ -Adic  $q$ -Integrals Representation on  $\mathbb{Z}_p$  Associated with the weighted  $q$ -Bernstein and  $q$ -Bernoulli polynomials, Journal of Inequalities and Applications, Article ID 513821, 8 pages, doi:10.1155/2011/513821.
- [22] C. S. Ryoo, A note on the weighted  $q$ -Euler numbers and polynomials, Adv. Stud. Contemp. Math. 21 (2011), 47-54.
- [23] H. Y. Lee, N. S. Jung, and C. S. Ryoo, Some Identities of the Twisted  $q$ -Genocchi Numbers and Polynomials with weight  $\alpha$  and  $q$ -Bernstein Polynomials with weight  $\alpha$ , Abstract and Applied Analysis, Volume 2011 (2011), Article ID 123483, 9 pages.
- [24] N. S. Jung, H. Y. Lee and C. S. Ryoo, Some Relations between Twisted  $(h, q)$ -Euler Numbers with Weight  $\alpha$  and  $q$ -Bernstein Polynomials with Weight  $\alpha$ , Discrete Dynamics in Nature and Society, Volume 2011 (2011), Article ID 176296, 11 pages.

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics,  
27310 Gaziantep, TURKEY  
*E-mail address:* mtsrkn@hotmail.com

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics,  
27310 Gaziantep, TURKEY  
*E-mail address:* acikgoz@gantep.edu.tr